

SEP 05 2007


Application No.: 10/664,671

Docket No.: JCLA12230-R2

AMENDMENTSIn the Claims:

Please amend the claims as follows:

1. (currently amended) A compound of formula (1):



wherein R^1 represents halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, $-\text{OCF}_3$, $-\text{OCF}_2\text{H}$, $-\text{N}=\text{C}=\text{O}$, $-\text{N}=\text{C}=\text{S}$, or alkyl having from 1 to 20 carbon atoms, and any $-\text{CH}_2-$ of the alkyl may be substituted with $-\text{O}-$, $-\text{S}-$, $-\text{CO}-$, $-\text{COO}-$, $-\text{OCO}-$, $-\text{CH}=\text{CH}-$, $-\text{CF}=\text{CF}-$ or $-\text{C}\equiv\text{C}-$, and any hydrogen thereof may be substituted with halogen or $-\text{CN}$; R^2 , R^3 and R^5 each independently represent hydrogen or alkyl having from 1 to 3 carbon atoms; A^1 , A^2 , A^3 and A^4 each independently represent 1,4-cyclohexylene, 1,4-cyclohexenylene, 1,4-phenylene, naphthalene-2,6-diyl, tetrahydronaphthalene-2,6-diyl, fluorene-2,7-diyl, bicyclo[2.2.2]octane-1,4-diyl or bicyclo[3.1.0]hexane-3,6-diyl, and in these rings, any $-\text{CH}_2-$ may be substituted with $-\text{O}-$, and any $-\text{CH}=\text{CH}-$ may be substituted with $-\text{N}=\text{N}-$, and in these rings, any hydrogen may be substituted with halogen or alkyl having from 1 to 5 carbon atoms; Z^1 , Z^2 and Z^3 each independently represent a single bond, $-(\text{CH}_2)_a-$, $-\text{O}(\text{CH}_2)_a-$, $-(\text{CH}_2)_a\text{O}-$, $-\text{O}(\text{CH}_2)_a\text{O}-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{COO}-$, $-\text{OCO}-$, $-(\text{CF}_2)_2-$, $-\text{C}\equiv\text{C}-\text{COO}-$, $-\text{OCO}-\text{C}\equiv\text{C}-$, $-\text{CH}=\text{CH}-(\text{CH}_2)_2-$, $-(\text{CH}_2)_2-\text{CH}=\text{CH}-$, $-\text{CF}=\text{CF}-$,

Application No.: 10/664,671

Docket No.: JCLA12230-R2

$-\text{C}\equiv\text{C}-\text{HC}=\text{CH}-$, or $-\text{CH}=\text{CH}-\text{C}\equiv\text{C}-$, and a indicates an integer of from 1 to 20; Z^4 represents a single bond or α,ω -alkylene having from 1 to 4 carbon atoms, and when Z^4 represents α,ω -alkylene having 3 or 4 carbon atoms, the $-\text{CH}_2-$ thereof directly bonded with A^4 may be substituted with $-\text{O}-$, $-\text{S}-$, $-\text{COO}-$ or $-\text{OCO}-$; m , n and q each independently indicates 0, 1 or 2, but $m+n+q\geq 1$;

wherein when $m+n+q=1$, any $-\text{CH}_2-$ of the alkyl represented by R^1 is not substituted with $-\text{CO}-$ and Z^4 is a single bond; ~~and~~

~~wherein~~ when $m+n+q=1$, Z^4 is a single bond and A^4 represents 1,4-phenylene, Z^1 , Z^2 and Z^3 each is not a single bond; and

when $m+n+q=2$, A^1 , A^2 , A^3 and A^4 are 1,4-phenylene and Z^1 , Z^2 , Z^3 and Z^4 are single bonds, any $-\text{CH}_2-$ of the alkyl represented by R^1 is not substituted with $-\text{CO}-$.

2. (original) A compound as claimed in claim 1, in which R^5 in formula (1) is hydrogen.

3. (original) A compound as claimed in claim 2, in which R^2 and R^3 in formula (1) in claim 1 are hydrogen.

4. (original) A compound as claimed in claim 3, in which A^1 , A^2 , A^3 and A^4 in formula (1) in claim 1 are independently any of 1,4-cyclohexylene or 1,4-phenylene, and any hydrogen in these rings may be substituted with halogen.





Application No.: 10/664,671

Docket No.: JCLA12230-R2

5. (previously presented) A compound as claimed in claim 3, in which A^1 , A^2 , A^3 and A^4 in formula (1) in claim 1 are independently any of 1,4-cyclohexylene or 1,4-phenylene, and any hydrogen in these rings may be substituted with halogen; and Z^1 , Z^2 and Z^3 are independently any of a single bond, $-(CH_2)_a-$, $-O(CH_2)_a-$, $-(CH_2)_aO-$, $-O(CH_2)_aO-$, $-CH=CH-$, $-C\equiv C-$, $-COO-$, or $-OCO-$.

6. (original) A compound as claimed in claim 5, in which Z^4 in formula (1) in claim 1 is a single bond.

7. (currently amended) Any one compound of formulae (a) to (d):



wherein R^1 represents halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, $-\text{OCF}_3$, $-\text{OCF}_2\text{H}$, $-\text{N}=\text{C}=\text{O}$, $-\text{N}=\text{C}=\text{S}$, or alkyl having from 1 to 20 carbon atoms, and any $-\text{CH}_2-$ of the alkyl may be substituted with $-\text{O}-$, $-\text{S}-$, $-\text{CO}-$, $-\text{COO}-$, $-\text{OCO}-$, $-\text{CH}=\text{CH}-$, $-\text{CF}=\text{CF}-$ or $-\text{C}\equiv\text{C}-$, and any hydrogen thereof may be substituted with halogen or $-\text{CN}$; A^1 , A^2 , A^3 and A^4 each independently

Application No.: 10/664,671

Docket No.: JCLA12230-R2

represent 1,4-cyclohexylene, 1,4-cyclohexenylene, 1,4-phenylene, naphthalene-2,6-diyl, tetrahydronaphthalene-2,6-diyl, fluorene-2,7-diyl, bicyclo[2.2.2]octane-1,4-diyl or bicyclo[3.1.0]hexane-3,6-diyl, and in these rings, any

$-\text{CH}_2-$ may be substituted with $-\text{O}-$, and any $-\text{CH}=\text{}$ may be substituted with $-\text{N}=\text{}$, and in these rings, any hydrogen may be substituted with halogen or alkyl having from 1 to 5 carbon atoms; Z^1 , Z^2 and Z^3 each independently represent a single bond, $-(\text{CH}_2)_a-$,

$-\text{O}(\text{CH}_2)_a-$, $-(\text{CH}_2)_a\text{O}-$, $-\text{O}(\text{CH}_2)_a\text{O}-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{COO}-$, $-\text{OCO}-$, $-(\text{CF}_2)_2-$,

$-\text{C}\equiv\text{C}-\text{COO}-$, $-\text{OCO}-\text{C}\equiv\text{C}-$, $-\text{CH}=\text{CH}-(\text{CH}_2)_2-$, $-(\text{CH}_2)_2-\text{CH}=\text{CH}-$, $-\text{CF}=\text{CF}-$,

$-\text{C}\equiv\text{C}-\text{HC}=\text{CH}-$, or $-\text{CH}=\text{CH}-\text{C}\equiv\text{C}-$, and a indicates an integer of from 1 to 20; Z^4 represents a single bond or α,ω -alkylene having from 1 to 4 carbon atoms, and when Z^4 represents α,ω -alkylene having 3 or 4 carbon atoms, the $-\text{CH}_2-$ thereof directly bonded with A^4 may be substituted with $-\text{O}-$, $-\text{S}-$, $-\text{COO}-$ or $-\text{OCO}-$, and

wherein in formula (a),

any $-\text{CH}_2-$ of the alkyl represented by R^1 is not substituted with $-\text{CO}-$,

Z^4 is a single bond, and

Z^1 is not a single bond when A^4 represents 1,4-phenylene; and

in formula (b),

when A^1 , A^2 and A^4 are 1,4-phenylene and Z^1 , Z^2 and Z^4 are single bonds, any

$-\text{CH}_2-$ of the alkyl represented by R^1 is not substituted with $-\text{CO}-$.

Application No.: 10/664,671

Docket No.: JCLA12230-R2

8. (previously presented) A compound as claimed in claim 7, in which R^1 in formulae (a) to (d) is halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, $-\text{OCF}_3$, $-\text{OCF}_2\text{H}$, alkyl having from 1 to 10 carbon atoms, alkoxy having from 1 to 10 carbon atoms, alkoxyalkyl having from 2 to 10 carbon atoms, or alkenyl having from 2 to 10 carbon atoms; A^1 , A^2 , A^3 and A^4 are independently any of 1,4-cyclohexylene or 1,4-phenylene, and in these rings, any hydrogen may be substituted with halogen; Z^1 , Z^2 and Z^3 are independently any of a single bond, $-(\text{CH}_2)_2-$, $-(\text{CH}_2)_4-$, $-\text{OCH}_2-$, $-\text{O}(\text{CH}_2)_3-$, $-\text{CH}_2\text{O}-$, $-(\text{CH}_2)_3\text{O}-$, $-\text{O}(\text{CH}_2)_2\text{O}-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{COO}-$, $-\text{OCO}-$, $-(\text{CF}_2)_2-$, or $-\text{CF}=\text{CF}-$; Z^4 is a single bond.

9. (previously presented) A liquid-crystal composition containing at least two polymerizable compounds, in which at least one polymerizable compound is the compound of claim 1.

10. (previously presented) A liquid-crystal composition, which contains at least two polymerizable compounds in which all the polymerizable compounds are the compounds of claim 1.


11. (previously presented) A liquid-crystal composition, which contains at least two polymerizable compounds that comprise at least one compound of claim 1 and at least one polymerizable compound except the compound.

Application No.: 10/664,671

Docket No.: JCLA12230-R2

12. (original) A liquid-crystal composition as claimed in claim 9, which additionally contains an optically-active compound.

13. (currently amended) A polymer having a constitutional unit of formula (2):



wherein R^1 represents hydrogen, halogen, $-\text{CN}$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, $-\text{OCF}_3$, $-\text{OCF}_2\text{H}$, $-\text{N}=\text{C}=\text{O}$, $-\text{N}=\text{C}=\text{S}$, or alkyl having from 1 to 20 carbon atoms, and any $-\text{CH}_2-$ of the alkyl may be substituted with $-\text{O}-$, $-\text{S}-$, $-\text{CO}-$, $-\text{COO}-$, $-\text{OCO}-$, $-\text{CH}=\text{CH}-$, $-\text{CF}=\text{CF}-$ or $-\text{C}\equiv\text{C}-$, and any hydrogen thereof may be substituted with halogen or $-\text{CN}$; R^2 , R^3 and R^5 each independently represent hydrogen or an alkyl having from 1 to 3 carbon atoms; A^1 , A^2 , A^3 and A^4 each independently represent 1,4-cyclohexylene, 1,4-cyclohexenylene, 1,4-phenylene, naphthalene-2,6-diyl, tetrahydronaphthalene-2,6-diyl, fluorene-2,7-diyl, bicyclo[2.2.2]octane-1,4-diyl or bicyclo[3.1.0]hexane-3,6-diyl, and in these rings, any $-\text{CH}_2-$ may be substituted with $-\text{O}-$, and any $-\text{CH}=\text{CH}-$ may be substituted with $-\text{N}=\text{N}-$, and in these rings, any hydrogen may be substituted with halogen or alkyl having from 1 to 5 carbon atoms; Z^1 , Z^2 and Z^3 each independently represent a single bond, $-(\text{CH}_2)_a-$, $-\text{O}(\text{CH}_2)_a-$, $-(\text{CH}_2)_a\text{O}-$, $-\text{O}(\text{CH}_2)_a\text{O}-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{COO}-$, $-\text{OCO}-$, $-(\text{CF}_2)_2-$, $-\text{C}\equiv\text{C}-\text{COO}-$, $-\text{OCO}-\text{C}\equiv\text{C}-$, $-\text{CH}=\text{CH}-(\text{CH}_2)_2-$, $-(\text{CH}_2)_2-\text{CH}=\text{CH}-$, $-\text{CF}=\text{CF}-$, $-\text{C}\equiv\text{C}-\text{HC}=\text{CH}-$, $-\text{CH}=\text{CH}-\text{C}\equiv\text{C}-$, $-\text{OCF}_2-$, or $-\text{CF}_2\text{O}-$, and a indicates an integer of from 1 to

Application No.: 10/664,671

Docket No.: JCLA12230-R2

20; Z^4 represents a single bond or α,ω -alkylene having from 1 to 4 carbon atoms, and any $-\text{CH}_2-$ of the alkylene may be substituted with $-\text{O}-$, $-\text{S}-$, $-\text{COO}-$ or $-\text{OCO}-$; and m , n and q each independently indicate 0, 1 or 2; and wherein when $m+n+q=1$, any $-\text{CH}_2-$ of the alkyl represented by R^1 is not substituted with $-\text{CO}-$ and Z^4 is a single bond; and when $n+m+q=2$, A^1 , A^2 , A^3 and A^4 are 1,4-phenylene and Z^1 , Z^2 , Z^3 and Z^4 are single bonds, any $-\text{CH}_2-$ of the alkyl represented by R^1 is not substituted with $-\text{CO}-$.

14. (original) A polymer as claimed in claim 13, in which R^5 in formula (2) is hydrogen.

15. (original) A polymer as claimed in claim 13, in which R^2 , R^3 and R^5 are hydrogen.

16. (original) A polymer as claimed in claim 13, in which R^2 , R^3 and R^5 are hydrogen; A^1 , A^2 , A^3 and A^4 are independently any of 1,4-cyclohexylene or 1,4-phenylene, and any hydrogen in these rings may be substituted with halogen.

17. (original) A polymer as claimed in claim 13, in which R^2 , R^3 and R^5 are hydrogen; A^1 , A^2 , A^3 and A^4 are independently any of 1,4-cyclohexylene or 1,4-phenylene, and any hydrogen in these rings may be substituted with halogen; and Z^1 , Z^2 and Z^3 are independently any of a single bond, $-(\text{CH}_2)_n-$, $-\text{O}(\text{CH}_2)_n-$, $-(\text{CH}_2)_n\text{O}-$, $-\text{O}(\text{CH}_2)_n\text{O}-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{COO}-$, $-\text{OCO}-$, $-\text{OCF}_2-$, or $-\text{CF}_2\text{O}-$.


Application No.: 10/664,671

Docket No.: JCLA12230-R2

18. (original) A polymer as claimed in claim 13, in which R^2 , R^3 and R^5 are hydrogen; A^1 , A^2 , A^3 and A^4 are independently any of 1,4-cyclohexylene or 1,4-phenylene, and any hydrogen in these rings may be substituted with halogen; Z^1 , Z^2 and Z^3 are independently any of a single bond, $-(CH_2)_a-$, $-O(CH_2)_a-$, $-(CH_2)_aO-$, $-O(CH_2)_aO-$, $-CH=CH-$, $-C\equiv C-$, $-COO-$, $-OCO-$, $-OCF_2-$, or $-CF_2O-$, and Z^4 is a single bond.

19. (original) A polymer as claimed in claim 13, in which R^1 in formula (2) is hydrogen, halogen, $-CN$, $-CF_3$, $-CF_2H$, $-CFH_2$, $-OCF_3$, $-OCF_2H$, alkyl having from 1 to 10 carbon atoms, alkoxy having from 1 to 10 carbon atoms, alkoxyalkyl having from 2 to 10 carbon atoms, or alkenyl having from 2 to 10 carbon atoms; R^2 , R^3 and R^5 are hydrogen; A^1 , A^2 , A^3 and A^4 are independently any of 1,4-cyclohexylene or 1,4-phenylene, and in these rings, any hydrogen may be substituted with halogen; Z^1 , Z^2 and Z^3 are independently any of a single bond, $-(CH_2)_2-$, $-(CH_2)_4-$, $-OCH_2-$, $-O(CH_2)_3-$, $-CH_2O-$, $-(CH_2)_3O-$, $-O(CH_2)_2O-$, $-CH=CH-$, $-C\equiv C-$, $-COO-$, $-OCO-$, $-(CF_2)_2-$, $-CF=CF-$, $-OCF_2-$ or $-CF_2O-$; Z^4 is a single bond.


20. (previously presented) A polymer that is obtained through homopolymerization of one compound of claim 1 and has a constitutional unit of formula (2):



Application No.: 10/664,671

Docket No.: JCLA12230-R2

21. (previously presented) A polymer that is obtained from the liquid-crystal composition of claim 9 and has a constitutional unit of formula (2):



22. (previously presented) An optically-anisotropic material of the polymer of claim 13.



23. (previously presented) A liquid-crystal display device, which contains the polymer of claim 13.

24. (original) A liquid-crystal display device, which contains the optically-anisotropic material of claim 22.

25. (currently amended) A method for producing a vinyl ketone compound of formula (1b), which comprises reacting one molar equivalent of a compound of formula (1a) with from 1 to 10 molar equivalents of a Lewis acid at -70°C to 200°C, followed by dehydrohalogenating the resulting compound:

Application No.: 10/664,671

Docket No.: JCLA12230-R2



wherein R^4 represents hydrogen, halogen, $-\text{OH}$, $-\text{CN}$, $-\text{CF}_3$, $-\text{CF}_2\text{H}$, $-\text{CFH}_2$, $-\text{OCF}_3$, $-\text{OCF}_2\text{H}$, $-\text{N}=\text{C}=\text{O}$, $-\text{N}=\text{C}=\text{S}$, or alkyl having from 1 to 20 carbon atoms, and any $-\text{CH}_2-$ of the alkyl may be substituted with $-\text{O}-$, $-\text{S}-$, $-\text{CO}-$, $-\text{COO}-$, $-\text{OCO}-$, $-\text{CH}=\text{CH}-$, $-\text{CF}=\text{CF}-$ or $-\text{C}\equiv\text{C}-$, and any hydrogen thereof may be substituted with halogen or $-\text{CN}$; R^2 , R^3 and R^5 each independently represent hydrogen or an alkyl having from 1 to 3 carbon atoms; A^1 , A^2 , A^3 and A^4 each independently represent 1,4-cyclohexylene, 1,4-cyclohexenylene, 1,4-phenylene, naphthalene-2,6-diyl, tetrahydronaphthalene-2,6-diyl, fluorene-2,7-diyl, bicyclo[2.2.2]octane-1,4-diyl or bicyclo[3.1.0]hexane-3,6-diyl, and in these rings, any $-\text{CH}_2-$ may be substituted with $-\text{O}-$, and any $-\text{CH}=\text{CH}-$ may be substituted with $-\text{N}=\text{N}-$, and in these rings, any hydrogen may be substituted with halogen or alkyl having from 1 to 5 carbon atoms; Z^1 , Z^2 and Z^3 each independently represent a single bond, $-(\text{CH}_2)_a-$, $-\text{O}(\text{CH}_2)_a-$, $-(\text{CH}_2)_a\text{O}-$, $-\text{O}(\text{CH}_2)_a\text{O}-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{COO}-$, $-\text{OCO}-$, $-(\text{CF}_2)_2-$, $-\text{C}\equiv\text{C}-\text{COO}-$, $-\text{OCO}-\text{C}\equiv\text{C}-$, $-\text{CH}=\text{CH}-(\text{CH}_2)_2-$, $-(\text{CH}_2)_2-\text{CH}=\text{CH}-$, $-\text{CF}=\text{CF}-$, $-\text{C}\equiv\text{C}-\text{HC}=\text{CH}-$, $-\text{CH}=\text{CH}-\text{C}\equiv\text{C}-$, $-\text{OCF}_2-$ or $-\text{CF}_2\text{O}-$, and a indicates an integer of from 1 to 20; Z^4 represents a single bond or α,ω -alkylene having from 1 to 4 carbon atoms, and

Application No.: 10/664,671

Docket No.: JCLA12230-R2

any $-\text{CH}_2-$ of the alkylene may be substituted with $-\text{O}-$, $-\text{S}-$, $-\text{COO}-$ or $-\text{OCO}-$; m, n and q each independently indicate 0, 1 or 2; Hal represents chlorine, bromine or iodine, wherein
when $m+n+q=1$, any $-\text{CH}_2-$ of the alkyl represented by R^1 is not substituted with $-\text{CO}-$ and Z^4 is a single bond; and
when $m+n+q=2$, A^1 , A^2 , A^3 and A^4 are 1,4-phenylene and Z^1 , Z^2 , Z^3 and Z^4 are single bonds,
any $-\text{CH}_2-$ of the alkyl represented by R^1 is not substituted with $-\text{CO}-$.